

Scotland's Rural College

## Relative emissions intensity of dairy production systems: employing different functional units in life-cycle assessment

Ross, SA; Topp, CFE; Ennos, RA; Chagunda, MGG

*Published in:*  
Animal

*DOI:*  
[10.1017/S1751731117000052](https://doi.org/10.1017/S1751731117000052)

First published: 10/02/2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

### *Citation for pulished version (APA):*

Ross, SA., Topp, CFE., Ennos, RA., & Chagunda, MGG. (2017). Relative emissions intensity of dairy production systems: employing different functional units in life-cycle assessment. *Animal*, 11(8), 1381 - 1388.  
<https://doi.org/10.1017/S1751731117000052>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Relative emissions intensity of dairy production systems: employing different functional units in life cycle assessment**

Stephen A. Ross <sup>1,a</sup>, Cairistiona F.E. Topp <sup>2</sup>, Richard A. Ennos <sup>3</sup>, Mizeck G.G. Chagunda <sup>1</sup>

<sup>1</sup> Future Farming Systems Group, SRUC, Edinburgh, UK

<sup>2</sup> Crop and Soil Systems Systems Group, SRUC, Edinburgh, UK

<sup>3</sup> School of Biological Sciences, University of Edinburgh, Edinburgh, UK

<sup>a</sup> Singapore University of Technology and Design, Singapore

Corresponding author: Mizeck Chagunda

E-mail: mizeck.chagunda@sruc.ac.uk

Short title: Functional units in life cycle assessment of dairy

## **Abstract**

This study aimed to assess the merit and suitability of individual functional units (FU) in expressing greenhouse gas emissions intensity in different dairy production systems. A FU provides a clearly defined and measurable reference to which input and output data are normalised. This enables the results from life cycle assessment (LCA) of different systems to be treated as functionally equivalent. Although the methodological framework of LCA has been standardised, selection of an appropriate FU remains ultimately at the discretion of the individual study. The aim of the present analysis was to examine the effect of different FU on the emissions intensities of different dairy production systems. Analysis was based on seven years of data (2004-2010) from four Holstein-Friesian dairy systems at Scotland's Rural College's long-term genetic and management systems project, the Langhill herd. Implementation of LCA accounted for the environmental impacts of the whole-farm systems and their production of milk from 'cradle to farm gate'. Emissions intensity was determined as kilograms of carbon dioxide equivalents referenced to six FU: United Kingdom livestock units, energy corrected milk yield, total combined milk solids yield, on-farm land used for production, total combined on- and off-farm land used for production, and the proposed new FU – energy corrected milk yield per hectare of total land used. Energy corrected milk was the FU most effective for reflecting differences between the systems. FU which incorporated a land-related aspect did not find difference between systems which were managed under the same forage regime, despite their comprising different genetic lines. Employing on-farm land as the FU favoured grazing systems. The proposed dual FU combining both productivity and land use did not differentiate between emissions intensity of systems as effectively as the productivity-based units. However, this dual unit

displayed potential to quantify in a simple way the positive or negative outcome of trade-offs between land and production efficiencies, in which improvement in emissions intensity using one FU may be accompanied by deterioration using another FU. The perceived environmental efficiencies of different dairy production systems in terms of their emissions intensities were susceptible to change based upon the FU employed, and hence the FU used in any study needs to be taken into account in the interpretation of results.

**Keywords:** Life Cycle Assessment, Functional Unit, Dairy Cow, Greenhouse Gas

## **Implications**

Dairy production systems are key contributors of greenhouse gas emissions. Emissions intensity is estimated by life cycle assessment and is influenced by different feeding and management systems and dairy cows. The perceived environmental efficiency of different dairy systems can change depending on the functional unit to which emissions are referenced. Results from studies comparing the emissions intensity of different dairy production systems should be considered in context of the functional unit used, in order to appraise them in an informed manner.

## Introduction

Life cycle assessment (LCA) has become a leading tool employed in agriculture for environmental impact and greenhouse gas (GHG) emissions accounting at the whole-systems level. Favoured for its flexibility, LCA enables an account to be made of all system inputs, processes and outputs within a specified boundary. In order to improve transparency and consistency amongst studies, the international standard ISO 14040 was established (ISO, 2006), stipulating requirements and recommendations for the LCA decision-making process. Further frameworks attempting to institute consistency in LCA at national and industry-specific levels have been developed, such as PAS 2050 in the United Kingdom (BSI, 2011). By convention, LCA results must be referenced to a functional unit (FU), providing a clearly defined and measurable reference to which input and output data are normalised (ISO, 2006). For the purposes of GHG measurement, the FU can either be a single item of product or a generally accepted sales quantity (BSI, 2011). The selection of an appropriate FU is crucial when assessing and interpreting environmental impacts, in which an impact category such as GHG emissions may be referenced to several different FU (Haas *et al.*, 2001).

In LCA of dairy production systems, the FU has most frequently been a unit of milk (Yan *et al.*, 2011; Baldini *et al.*, 2016), the principal unit of production in dairy. Milk yields have commonly been corrected to standardised levels of milk fat and protein content, as milk composition commonly determines the value of raw milk to the processor. This enables better comparison between farms with different breeds or different feeding regimes (IDF, 2010). Other production-based FU have been employed, including the combined mass of milk fat and protein as milk solids (MS),

or simply as uncorrected raw milk yields (Baldini *et al.*, 2016). When the LCA boundary included post-farm processing of dairy products, results have been referenced to a unit of consumer product, such as processed packaged milk (Hospido *et al.*, 2003). Expressing emissions intensity per unit of productivity is essentially a ratio of undesirable versus desirable system outputs. Land use is also commonly used and satisfies a more conventional definition of efficiency, being a measure of system output versus system input. As a FU, land use can include the productive on-farm land required for grazing and forage production or the entire farm area. Recently, an area-based FU has also incorporated off-farm land required for production of purchased feeds (O'Brien *et al.*, 2012). The livestock themselves, expressed as the total number of livestock units (LSU) on the farm, has also been used as an FU (Haas *et al.*, 2001). Although concerted efforts have been made to establish consistency in the application of LCA to dairy production systems, selection of FU ultimately remains at the discretion of individual investigators.

Studies directly examining the effect of varying the FU on the results of LCA of different dairy production systems are sparse. Several studies have employed multiple FUs, therefore indirectly providing some assessment, albeit not necessarily as the primary aim (e.g. Haas *et al.*, 2001; O'Brien *et al.*, 2012). Results of these studies indicate that the perceived relative environmental efficiency of dairy production systems can change based on the FU employed. The objective of our study was to examine the effect of employing different FU on the estimated environmental efficiency of different dairy production systems and hence the ability to classify different production systems based on GHG emissions intensity.

## **Materials and Methods**

### *Dairy production systems and life cycle assessment*

The study was based on data from Scotland's Rural College's (SRUC) long-term genetic and management systems project, the Langhill herd, located at the SRUC Dairy Research Centre, Crichton Royal Farm, Dumfries, Scotland. Data used were collected from January 2004 to December 2010 from four distinct systems within a conventional farm. Holstein-Friesian cows were maintained in two feeding groups: high forage (HF) and low forage (LF). The HF systems aimed to provide 75% by dry matter of the herd's total mixed ration (TMR) diet from home-grown crops (i.e. ryegrass silage, whole-crop maize, wheat alkalage) and 25% from purchased concentrated feeds (e.g. distillers grains, rapeseed meal). Cows in the HF systems were also put outside to graze pasture, when available. In contrast, cows in the LF systems were fully housed and fed a diet of approximately 45% of the same home-grown forages and 55% purchased concentrated feeds. Within each forage system, animals comprised two contrasting genetic lines. Control (C) animals were bred to be of average United Kingdom (UK) genetic potential for milk fat and protein production, while Select (S) animals represented the top 5% of UK genetic potential. Maintaining the specific characteristics of these groups in a long-term genotype × feeding regime project resulted in four divergent dairy production systems: HFC, HFS, LFC and LFS. Cows were milked three times daily and received equal health and fertility treatments, and herd size was maintained at approximately 50 cows per system. C and S cows were managed together, and groups were kept in the same building when housed. All rations were formulated using the same preserved forages, and all young stock were managed together (Chagunda *et al.*, 2009).

Annual GHG emissions were estimated using LCA, covering all stages of dairy production from the extraction or acquisition of raw materials up to the point at which the product milk left the farm. Briefly, high-resolution data were collected for on-farm processes, including herd dynamics, electricity and fuel use, crop production, fertiliser application, as well as animals' diet formulation, feed intake and productivity. Impacts were assessed by applying Intergovernmental Panel on Climate Change (IPCC) Tier 2 methodology (IPCC, 2006). IPCC coefficients were used to estimate direct emissions, volatilization and leaching associated with application and storage of fertilizer and manure, and from crop residues (IPCC, 2006). UK emissions factors were used for production and transport of fertilisers, concentrated feeds and animal bedding (Carbon Trust, 2010), and for electricity and fuel use (DEFRA, 2011). System-specific Tier 3 emissions factors were estimated for enteric methane, manure methane and excreted nitrogen (Ross *et al.*, 2014). Life cycle emissions were allocated between two system outputs - the products milk and meat - using mass allocation. Emissions intensity was defined as the estimated global warming potential (GWP) of each system in kg of carbon dioxide equivalents (kg CO<sub>2</sub>e) per FU. Additional in-depth details of the LCA performed are described in Ross *et al.* (2014).

#### *Functional units*

The estimated annual GWP of dairy production systems was referenced to five FU employed in previous LCA studies of dairy production systems and to one further FU which incorporated a measure of both the productivity and land use of the systems (Table 1).



*Livestock units.* At system level, the LCA included not only milking cows but also all other age classes of livestock. Employing LSU enabled the entire herd to be included in the FU, applying a conversion factor for the relative ages and persistence of young and replacement animals. These were defined for calves aged 0-12 months, heifers 12-24 months and heifers over 24 months as 0.34, 0.65 and 0.80, respectively (DEFRA, 2010). Milking cows in different systems were corrected for average bodyweight and milk yield (ADAS, 1983), with values of 1.48, 1.64, 1.26 and 1.37 for LFC, LFS, HFC and HFS, respectively. All male calves were sold after weaning, and there were no bulls on the farm. Total annual populations of livestock were multiplied by their respective coefficients and summed to give an average population in LSU for each system.

*Energy corrected milk.* Milk yields amongst the production systems varied in energy content, which is defined by fat and protein contents (Table 2), potentially leading to differences in calculations if only the mass or volume of milk was considered. We used the equation of Sjaunja *et al.* (1990), commonly employed in LCA, to calculate energy corrected milk (ECM) with 35.0 g/kg milk fat and 32.0 g/kg protein, as follows:  $ECM (kg) = 0.25M + 12.2F + 7.7P$ , where M = annual milk yield (kg), F = milk fat content (kg), P = protein content (kg).

*Milk solids.* Data on milk yield were recorded for individual cows after each milking session. Milk fat and protein contents were recorded from samples collected from each cow, three times daily on one day each week. An estimate was thus made for the total mass of milk fat and protein yielded by each cow per calendar year.

Individual cow data were combined to give the total annual output of MS for each system.

*On-farm land-use.* On-farm land-use to provide forage crops (ryegrass silage, whole-crop wheat, maize) to each system was estimated based on the forage requirements of the system. Quantities of ensiled forages used were weighed during formulation of the daily TMR, and daily feed intake of cows was recorded using automated Hoko feeding gates (Insentec BV, Marknesse, NL). Forage crop requirements of each system were then related to harvested forage yields and estimated annual forage land requirements. Dry matter losses were considered during harvesting, ensiling and unloading of forages (Bastiman and Altman, 1985; MacDonald *et al.*, 1991). Land required by HF systems for pasture was similarly estimated based upon the predicted grazing dry matter intake (DMI) of cows and the available herbage per ha (Table 3). Applying the estimates from Bell *et al.* (2010), DMI of HF cows grazing pasture was 19.2 kg/day for C cows and 20.8 kg/day for S cows.

*Off-farm land-use.* Off-farm land associated with the external production of purchased feed components and bedding was estimated using the method of Bell *et al.* (2011) (Table 4). Firstly, total annual purchased feed required was estimated from recorded Hoko data and from TMR formulations, breaking the purchased fraction of the diets down into component ingredients. Land-use values for domestically produced purchased feed components (wheat, rapeseed, barley) were estimated using national data on crop yields (Craig & Logan, 2012; Scottish Government, 2012). Land-use for the internationally imported soya bean meal was sourced from the LCA food database (Nielsen *et al.*, 2007). Finally, allocations between co-

products of purchased feed components were made using mass-based factors from Cederberg and Mattsson (2000).

*Dual functional unit.* In this study we introduce a new FU including both the productivity and land use of systems. The estimated annual GWP of the dairy production system was divided by the total annual ECM yield, and this quotient further divided by the total on- and off-farm land use of the system for each year of the study. This dual FU thus incorporated the ratio of undesirable output (GHG) to desirable output (milk) per unit input (area of land) and therefore adhered to a more standardised output/input measure of efficiency.

#### *Statistical analysis*

Two statistical procedures were employed in the analysis. The effect of employing different FU upon the estimated environmental efficiency of dairy production systems was assessed using analysis of variance (ANOVA) employing the following generalised linear model (GLM),  $y_{ij} = \mu + S_i + Y_i + \varepsilon_{ij}$  where  $y_i$  was the total GWP of the dairy production system per FU (LSU, ECM, MS,  $Land_{farm}$ ,  $Land_{total}$ ,  $ECM/Land_{total}$ ),  $\mu$  was the overall mean,  $S_i$  was the fixed effect of dairy production system (LFC, LFS, HFC, HFS),  $Y_i$  was the random effect of calendar year (2004-2010), and  $\varepsilon_{ij}$  was the random error term. Fisher tests were used to assess the level of significance of contributing effects, and differences between dairy production systems were determined by conducting pairwise comparisons using the Tukey method.

The ability of different FU to classify different production systems' relative environmental efficiency was assessed using rank analysis. Using year as a repeated measure, systems were assigned a rank value from 1 to 4 in order of their relative emissions intensity, with rank 1 having the lowest GWP per FU and thus most efficient system. Rank analysis was performed using the Kruskal-Wallis test, a non-parametric equivalent to ANOVA. Significant differences between any two systems were assessed using the Mann-Whitney-Wilcoxon rank sum test. The rank analysis was repeated when referencing the GWP of systems to each of the six FU employed in this study. All statistical analyses were conducted using Minitab 16.

## **Results**

### *Emissions intensity*

The effect of the dairy production system on the overall GWP per FU was significantly different ( $P<0.001$ ) for each of the six FU. ECM was the only FU for which the GWPs of all four systems were significantly different from each other ( $P<0.001$ ) (Table 5). LFS had the lowest GWP per kg ECM, followed by LFC, HFS and HFC. Using MS as the FU, LFS again had the lowest GWP and HFC the highest. There was no significant difference between LFC and HFS per kg of MS. Using LSU as the FU, LFC was the most efficient, although not significantly different from LFS. LSU was the only FU which did not find a significant difference between LFS and HFC. Using  $\text{Land}_{\text{farm}}$  as the FU, GWP per ha of both HF systems was lower than those of both LF systems. Conversely, including off-farm land use ( $\text{Land}_{\text{total}}$ ), both LF systems had lower GWP per ha than both HF systems. Using the dual FU, which incorporated both productivity and land use, both LF systems were more

efficient than both HF systems. However, none of the three FU which incorporated land use found a significant difference between HFC and HFS or between LFC and LFS. Overall, LFS was the most efficient system for four of the six FU (ECM, MS, Land<sub>total</sub>, ECM/Land<sub>total</sub>).

### *Rank analysis*

For each of the six FU, median rankings of environmental efficiency of dairy production systems were significantly different ( $P<0.05$ ) (Table 6). The median rankings broadly reflected the relative order of system efficiency observed from ANOVA results, with two differences. Using Land<sub>total</sub> as the FU, HFS was the lowest ranked system and significantly different from the 3<sup>rd</sup> ranked HFC. Furthermore, the two most efficient systems when using the dual FU, LFS and LFC, were significantly different from each other according to the Mann-Whitney-Wilcoxon rank sum test, with LFS having the lower GWP. However, when using LSU or Land<sub>farm</sub> as the FU, LFS was ranked 4<sup>th</sup>.

### **Discussion**

LSU are commonly used in the dairy industry to compare stocking densities and nutritional requirements of animals; however, LSU have been infrequently used when interpreting outputs of LCA studies. When using LSU as the FU in this study, the LF systems had the lowest emissions intensity. When Haas *et al.* (2001) referenced emissions to LSU (defined as 500 kg of cow body weight), extensive grazing-based systems had lower emissions than intensive systems. This trend was not reflected in our results. Given the higher gross emissions of the LF regime (Ross *et al.*, 2014) and that systems had similar numbers of milking cows, the difference

observed was likely due to greater animal performance under LF. The UK LSU is based on a standard of 48,000 MJ of metabolisable energy, defined as “the feed energy allowance of a 625 kg Friesian cow and the production of a 40 kg calf, and 4,500 litres of milk at 3.6% butterfat and 8.6% solids-not-fat” (ADAS, 1983; DEFRA, 2010). Adjusting the number of LSU in each system based on corrections for liveweight and productivity stipulated by ADAS (1983), differences in observed cow performance amongst systems were embedded in the FU. From rank analysis, the emissions intensity of LFC was lower than that of the higher yielding LFS system. However, S genetic line animals had a higher feed and metabolisable energy intake and greater milk yield than those of the C line. This led to the S line having higher enteric methane emissions per cow and higher gross emissions associated with both forage and purchased feeds. Unlike using ECM as the FU for emissions intensity, the higher gross emissions associated with the S line were not sufficiently offset by the higher productivity when using LSU.

Milk yield corrected for milk fat and protein content is the most commonly applied FU in LCA studies of dairy production. Employing ECM incorporated the effect of disparity in milk production amongst systems when examining the GWP. This was also the only FU to find the emissions intensity of all four Langhill systems to be significantly different from each other. In a recent study in Ireland, O'Brien *et al.* (2012) found that a confinement system had higher emissions intensity than a grazing system per kg ECM. Our results disagree with these findings, but there were several differences in farm-management practices and methods between the studies. The Irish systems had substantially lower milk yield than the Langhill systems, and a further key difference was that 70-80% of concentrated feed

components of the Irish diets were internationally imported. The latter included maize from the USA and rapeseed meal and molasses from Germany, whereas in the Langhill systems, forage crops were grown on-farm, and the purchased components of the TMR were largely sourced from the same country. It is important to recognize that results of studies, in particular those of HF and LF systems in our study, depend on site-specific conditions, such as the purchase of certain concentrated feeds or the ability to grow certain crops locally. Langhill rations sourced a high percentage of their concentrated feeds from by-product grain from distilling and brewing industries, which has a considerably lower GHG emissions intensity than palm oil or soya bean meal imported from South America (Carbon Trust, 2010). Countries or regions differ considerably in their management preferences, climatic conditions, soil types, and availability or feasibility of crops. Brockman and Wilkins (2003) described variation amongst grass species in growth patterns, nutritional content, response to nitrogen and climate, and most importantly, yields. In turn, these factors influence the range of animal breeds and management systems available to dairy farmers, as well as the environmental impacts associated with them. This point highlights the importance of examining site-specific methods and farm-management practices when comparing LCA studies which use the same FU.

MS are another unit commonly used in the dairy industry, for example to compare the biological or production efficiencies of cows, but seldom used in LCA. This is likely because most dairy LCA studies draw their boundaries at the farm gate, at which the principal product is liquid milk. Exceptions to this have occurred in studies from countries where output of alternative dairy products such as dried milk, whey powder and butter exceeds that of liquid milk, such as New Zealand and Ireland.

This again is pertinent to the examination of site-specific practices. For studies interested in the life cycle of dairy-derived products such as yoghurt or cheese, referencing the GWP to MS may also be more appropriate. When using units which included dairy production (ECM and MS), results followed a similar trend, with LFS having the lowest emissions intensity and HFC the highest. However, unlike using ECM, there was no difference observed in GWP between LFC and HFS when using MS. This was likely due to a confounding effect introduced by the interaction of forage regime and genetic line in the production systems. Both fat and protein contents of milk were higher in HF systems, while milk yields were higher in the LF systems. This was not unexpected, as cows on a LF regime have historically been subject to milk-fat depression (Bauman and Griinari, 2001). Both milk yield and MS were higher with the S genetic line, consistent with its genetic potential. Thus while the LFC system yielded more raw milk containing fewer MS, the HFS produced less milk containing more MS. Although calculating ECM corrected for differences amongst systems' milk fat and protein contents, the impact of this confounding effect upon GWP was more pronounced when MS was the FU. This emphasises the importance of considering not only the GWP results, but also what information the FU may contain or tell about the dairy production systems examined.

Using land use as the FU satisfies the conventional definition of efficiency as a measure of system output to system input. Further, land area conforms to the ISO 14040 stipulation that the FU be a clearly defined and measurable reference to which input and output data are normalised (ISO, 2006). For studies which intend to inform national GHG inventory reporting, it is also necessary to choose a FU coupled with land for area-based processes (IPCC, 2006). In agriculture, this may include



emissions associated with applied fertilisers and crop production and residues, but it is not a requirement for animal emissions such as enteric CH<sub>4</sub>. The LCA process may include within its boundary emissions sources beyond those required for national inventory reporting; thus, an area-based unit is not a specific requirement of LCA. LCA studies have variously referenced GWP to on-farm pasture, combined on-farm pasture and forage-crop land, or to total on- and off-farm land, including that required to produce imported concentrated feeds and bedding (Haas *et al.*, 2001; van der Werf *et al.*, 2009; O'Brien *et al.*, 2012). In our study, on-farm land was the only FU for which both HF systems were more efficient than both LF systems. High overall GWP combined with lower on-farm land use due to the absence of grazing meant that LF systems had higher GWP per ha of on-farm land. In HF systems, higher on-farm land use resulted in lower GWP per ha. From an LCA point of view, including only on-farm area is not appropriate for examining results since arable land used to produce purchased feed, which could be located worldwide, will also be responsible for environmental impacts in the life cycle of milk production (Yan *et al.*, 2011). Thus, the GWP reported for conventional dairy production systems are often inexorably linked to the emissions intensities of crop production abroad.

As noted earlier, a FU serves as a measurable reference to which input and output data are normalised. In the same way that milk yields can be adjusted according to their fat and protein contents to ensure functional equivalence, should productive areas of land be adjusted in LCA to normalise their productivities across different locations? In doing this, GWP per ha would no longer reflect differences in crop yields, but only differences in downstream system specific factors such as animal genetics, the formulation of rations, and other farm-management factors. Noting very different areas of land used to produce equivalent quantities of grain, Audsley *et al.*

(1997) equated land areas of different systems by assuming that the difference in land used was managed as set-aside land (Audsley *et al.*, 1997). Wackernagel and Rees (1996) introduced the notion of a 'global hectare' with their concept of the Ecological Footprint. To equate land around the world, they quantified demand on biological resources by expressing all components of an impact as an equivalent area of land and sea with world average productivity. Yield and equivalence factors were used to convert actual physical areas from local ha to global ha. Scaled down to the level of agricultural systems, yield factors, obtained by dividing the local yield of a biological product by its global average yield (Wiedmann and Lenzen, 2007), can account for differences in productivity of a given crop and land type amongst countries.

Increasing efficiency of livestock production by improving genetic potential has been identified as a promising approach for reducing global emissions from livestock systems (Steinfeld *et al.*, 2006). In the Langhill systems, S animals were bred to be in the top 5% of UK genetic potential for milk fat and protein contents. Previous studies have noted lower emissions intensity of S systems for both enteric methane (Chagunda *et al.*, 2009) and overall GWP per unit ECM (Ross *et al.*, 2014). When using an area-based FU, however, results did not differ between C and S genetic lines managed under the same forage regime. S animals consume more feed than C animals and consequently require more land and crops to produce their feed. Thus, as milk production, emissions and land use increase, using an area-based FU does not reflect the specific improvements made by the Langhill selection criteria. This is not so much a problem of the FU as it is reflection of what the FU reveals about the systems. The total land FU is a valid measure of agricultural or emissions intensity,

and, in the event of no significant difference between systems under study, one could look to production or other criteria to evaluate systems.

Some LCA studies of dairy production have used a FU unique to the study, such as “1000 euros of gross farm income” (van der Werf *et al.*, 2009). As the selection of a FU often remains at the discretion of researchers, they have the opportunity to innovate new and diverse FU to provide a balanced or new perspective. These FU could incorporate a socio-economic aspect such as the monetary unit employed by van der Werf *et al.* (2009). The use of either ECM or MS as the FU, rather than raw milk yields, enables a study to account not just for milk quantity but quality, which in turn can translate into milk price. Considering the observed differences in emissions intensity between HF and LF systems, a farm-management decision to implement a given system, made for economic reasons, inexorably influences the system’s environmental impacts. Feed costs can account for around 80% of total variable costs of milk production (Shalloo *et al.*, 2004) and around 35% of the environmental impacts of the Langhill systems in this study (Ross *et al.*, 2014). In future, LCA should perhaps consider a FU which can reflect economic functions of a system, such as “GWP per 1000 euros of gross farm income” (van der Werf *et al.*, 2009) in addition to the production or land use. Alternatively, a FU could reflect competition for resources (e.g., per kg of non-human-edible feed) or reflect efficiency (e.g., per MJ of potential energy of system inputs such as fuel, feed). However, the perceived performance of any production system as a function of efficiency may have to do with the different ways in which efficiency is understood. For example, efficiency in an animal production system is commonly defined by its feed conversion ratio. This also raises the question of whether FU selection should be tailored to the intention of a study or use a widely employed, standard FU. Further, a FU in LCA should be

relevant to reference a wide range of impact categories. Haas *et al.* (2001) stated that environmental impacts at a regional or local level, such as nitrate or phosphate pollution of a lake, have a strong area-related aspect and must be minimised irrespective of production levels that farmers can achieve. Given the multifunctionality of agricultural systems, LCA results should be evaluated using both production and land-based FU to provide a balanced assessment.

The new FU developed in our study aimed to account for both the productivity and the total on- and off-farm land use of a system. Although LFS had significantly lower GWP per kg ECM, when using  $\text{ECM}/\text{Land}_{\text{total}}$  as the FU, there was no significant difference between the emissions intensity of LFC and LFS. Indeed, much like the two FU based on land use, there was no significant difference between systems under the same forage regime. Given the range of milk yields in the Langhill systems, it was perhaps surprising that using the dual FU was able to differentiate only between the emissions intensities of feeding regimes but not those of genetic lines. Despite incorporating milk yield, the using this FU did not reflect the difference amongst systems' GWP that the existing FU incorporating productivity were able to determine. However, when analysing the systems' relative efficiency rankings in our study, the dual FU did find LFS to have significantly lower GWP than LFC.

It has been noted in the literature that there is a lack of significant correlation between GWP per unit product and GWP per unit land (Casey and Holden 2005). Yan *et al.* (2011) defined how this correlation could be achieved with an equation incorporating the milk yield per cow, the stocking rate and ratio of on-farm to off-farm land use. We propose that using the dual FU can measure the outcome of trade-offs, for example an improvement in emissions intensity using ECM as the FU being accompanied by a deterioration when using  $\text{Land}_{\text{total}}$ , when assessing conversion

from one production system to another. In the present study, a decrease in GWP per the dual FU would be observed in either a win-win scenario, in which both measures improved, or a trade-off with positive outcome, in which GWP per the dual FU would decrease despite an increase in GWP per ha, for example. Thus the remaining scenarios, trade-off with negative outcome and lose-lose, would be reflected by an increase in GWP per the dual FU. In the context of the present study, a conversion from HFC or HFS to either LFC or LFS represent win-win scenarios, with a decrease in GWP per ECM, per Land<sub>total</sub> and the dual FU. A conversion from HFC to HFS represents a positive outcome, with a decrease in GWP per dual FU despite an increase in GWP per Land<sub>total</sub>. Godfray *et al.* (2010) stated that there is a pressing need in modern global agriculture for 'sustainable intensification', in which yields are increased without adverse environmental impact and without cultivating more land. To comply with this definition, it would be advantageous for LCA of dairy production systems to use a FU which could account for environmental impacts relative to yield and land use simultaneously. In our study, the dual FU yielded results broadly consistent with those using ECM or total land-use as the FU, and helps to assess the likelihood of a positive trade-off between GHG emissions and efficiencies of production and land use. After investigating a similar concept of trade-offs between GWP per ha and ECM per ha, the study of Hayashi *et al.* (2013) recommended that both expressions should be used complementarily. It is advisable that, for an appropriately balanced assessment, LCA should consider the underlying reasons behind trade-offs, and thus also evaluate emissions intensity per both land-oriented and product-oriented FU individually.

## Conclusions

The relative emissions intensity of different dairy production systems sometimes changed based upon the FU employed. Energy corrected milk was the FU most effective for reflecting differences between the systems. FUs which incorporated a land-related aspect found no difference between systems which were managed under the same forage regime, despite their comprising different genetic lines with considerably different productivity.

Results from LCA studies comparing dairy production systems should be considered in the context of the FU used, in order to appraise them in an informed manner. ECM yields should remain the primary FU for comparing impacts of dairy production systems, however, both a land-use-based FU and the dual FU should be used in addition, in order to evaluate trade-offs and present a balanced assessment.

## **Acknowledgements**

The authors are grateful to the farm and technical staff at Crichton Royal Farm, from where the data used in this study were obtained. We are also grateful for the contribution made by Gillian Reid at the SAC Rural Business Unit. This study was funded by the Scottish Government.

## **References**

- ADAS 1983. Livestock Units Handbook. Agricultural Development Advisory Service. Booklet 2267. Ministry of Agriculture, Fisheries and Food, London, UK.
- Audsley E, Alber S, Clift R, Cowell S, Crettaz P, Gaillard G, Hausheer J, Jolliett O, Klejn R, Mortensen B, Pearce D, Roger E, Teulon H, Weidema B and van Zeijts H 1997. Harmonisation of environmental life cycle assessment for agriculture. Final Report,

507           Concerted Action AIR3-CT94-2028, European Commission, DG VI Agriculture,  
508           Brussels.

509   Baldini C, Gardoni D and Guarino M 2016. A critical review of the recent evolution of Life  
510           Cycle Assessment applied to milk production. *Journal of Cleaner Production*. Article in  
511           press. <http://dx.doi.org/10.1016/j.jclepro.2016.06.078>

512   Bastiman G and Altman JBF 1985. Losses at various stages of silage making. *Research and*  
513           *Development in Agriculture* 2, 19-25.

514   Bauman DE and Griinari JM 2001. Regulation and nutritional manipulation of milk fat: low-fat  
515           milk syndrome. *Livestock Production Science* 70, 15-29.

516   Bell MJ, Wall E, Russell G, Morgan C and Simm G 2010. Effect of breeding for milk yield,  
517           diet and management on enteric methane emissions from dairy cows. *Animal*  
518           *Production Science* 50, 817-826.

519   Bell MJ, Wall E, Russell G, Simm G and Stott AW 2011. The effect of improving cow  
520           productivity, fertility, and longevity on the global warming potential of dairy systems.  
521           *Journal of Dairy Science* 94, 3662-3678.

522   Brockman JS and Wilkins RJ 2003. Grassland. In: Soffe RJ (ed) Primrose McConnell's *The*  
523           *Agricultural Notebook*, 20<sup>th</sup> Edition. Oxford, Wiley-Blackwell, 131-176.

524   BSI 2011. PAS 2050:2011 Specification for the assessment of the life cycle greenhouse gas  
525           emissions of goods and services. British Standards Institution, London, UK.

526   Carbon Trust 2010. Footprint Expert Feed Database version 1.1. Carbon Trust Footprinting  
527           Company.

528   Casey JW and Holden NM 2005. The relationship between greenhouse gas emissions and  
529           the intensity of milk production in Ireland. *Journal of Environmental Quality* 34, 429-  
530           436.

531   Cederberg C and Mattsson B 2000. Life cycle assessment of milk production - a comparison  
532           of conventional and organic farming. *Journal of Cleaner Production* 8, 49-60.

533 Chagunda MGG, Römer DAM, and Roberts DJ 2009. Effect of genotype and feeding regime  
534 on enteric methane, non-milk nitrogen and performance of dairy cows during the winter  
535 feeding period. *Livestock Science* 122, 323-332.

536 Craig K, and Logan R 2012. *The Farm Management Handbook 2012/13 – The UK reference*  
537 *for farm business management*. 33<sup>rd</sup> Edition. SAC Farm and Rural Business Services,  
538 Edinburgh, UK.

539 DEFRA 2010. *Definitions of terms used in farm business management (3rd edition)*. The  
540 Department for Environment, Food and Rural Affairs, London, UK.

541 DEFRA 2011. *Guidelines to DEFRA / DECC's GHG conversion factors for company*  
542 *reporting*. The Department of Energy and Climate Change (DECC) and the  
543 Department for Environment, Food and Rural Affairs (DEFRA).

544 Godfray HCJ, Beddington JR, Crute IR, Haddad L, Lawrence D, Muir JF, Pretty J, Robinson  
545 S, Thomas SM and Toulmin C 2010. Food security: The challenge of feeding 9 billion  
546 people. *Science* 327, 812-818.

547 Haas G, Wetterich F, and Köpke U 2001. Comparing intensive, extensified and organic  
548 grassland farming in southern Germany by process life cycle assessment. *Agriculture*  
549 *Ecosystems & Environment* 83, 43-53.

550 Hayashi K 2013. Practical recommendations for supporting agricultural decisions through life  
551 cycle assessment based on two alternative views of crop production: the example of  
552 organic conversion. *International Journal of Life Cycle Assessment* 18, 331-339.

553 Hospido A, Moreira MT and Feijoo G 2003. Simplified life cycle assessment of Galician milk  
554 production. *International Dairy Journal* 13, 783-796.

555 IDF 2010. *A common carbon footprint approach for dairy: the IDF guide to a standard life*  
556 *cycle assessment methodology for the dairy sector*. Bulletin of the International Dairy  
557 Federation 445/2010. Brussels, Belgium

558 IPCC 2006. *2006 IPCC Guidelines for National Greenhouse Gas Inventories*, Prepared by  
559 the National Greenhouse Gas Inventories Programme. Eggleston H.S., Buendia L.,  
560 Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Hayama, Japan.



561 ISO 2006. International Standard 14044:2006. Environmental management - Life Cycle  
562 Assessment: requirements and guidelines. International Standards Organisation,  
563 Geneva, Switzerland.

564 MacDonald P, Henderson AR and Heron SJE 1991. The Biochemistry of Silage. 2nd edition.  
565 Chalcombe Publications, Marlow, UK.

566 Nielsen PH, Neilsen AM, Weidema BP, Dalgaard R and Halberg N 2007. LCA food  
567 database. Danish Institute of Agricultural Sciences, Aarhus, Denmark. [http://](http://www.lcafood.dk/database)  
568 [www.lcafood.dk/database](http://www.lcafood.dk/database).

569 O'Brien D, Shalloo L, Patton J, Buckley F, Grainger C, and Wallace M 2012. A life cycle  
570 assessment of seasonal grass-based and confinement dairy farms. Agricultural  
571 Systems 107, 33-46.

572 Ross SA, Chagunda MGG, Topp CFE and Ennos R 2014. Effect of cattle genotype and  
573 feeding regime on greenhouse gas emissions intensity in high producing dairy cows.  
574 Livestock Science 170, 158-171.

575 Rural Business Unit 2009. The SAC carbon calculator and carbon management services –  
576 capability statement. SAC Rural Business Unit, Penicuik, UK.

577 Scottish Government 2012. Abstract of Scottish agricultural statistics 1982-2012. The  
578 Scottish Government, Edinburgh, UK.  
579 [http://www.gov.scot/Topics/Statistics/Browse/Agriculture-](http://www.gov.scot/Topics/Statistics/Browse/Agriculture-Fisheries/PubAbstract/Abstract2012)  
580 [Fisheries/PubAbstract/Abstract2012](http://www.gov.scot/Topics/Statistics/Browse/Agriculture-Fisheries/PubAbstract/Abstract2012)

581 Sjaunja LO, Baevre L, Junkkarinen L, Pedersen J, and Setälä J 1990. A Nordic proposal for  
582 an energy corrected milk (ECM) formula. 27th Session of the International Commission  
583 for Breeding and Productivity of Milk Animals, Paris, France.

584 Shalloo L, Kennedy J, Wallace M, Rath M and Dillon P 2004. The economic impact of cow  
585 genetic potential for milk production and concentrate supplementation level on the  
586 profitability of pasture based systems under different EU milk quota scenarios. Journal  
587 of Agricultural Science. 142, 357-369.

588 van der Werf HMG, Kanyarushoki C and Corson MS 2009. An operational method for the  
589 evaluation of resource use and environmental impacts of dairy farms by life cycle  
590 assessment. *Journal of Environmental Management* 90, 3643-3652.

591 Wackernagel M and Rees W 1996. *Our Ecological Footprint: Reducing human impact on the*  
592 *Earth*. Gabriola Island, BC, Canada: New Society Publishers.

593 Wiedmann T and Lenzen M 2007. On the conversion between local and global hectares in  
594 Ecological Footprint analysis. *Ecological Economics* 60, 673-677.

595 Yan M, Humphreys J and Holden NM 2011. An evaluation of life cycle assessment of  
596 European milk production. *Journal of Environmental Management* 92, 372-379.

597

598 **Tables**

599

600 **Table 1.** Functional units employed in the life cycle assessment

Functional Unit	Abbreviation	Unit
United Kingdom livestock unit	LSU	n
Energy corrected milk yield	ECM	kg
Total combined milk solids yield	MS	kg
On-farm land used for production	Land <sub>farm</sub>	ha
Combined on- and off-farm land used for production	Land <sub>total</sub>	ha
Energy corrected milk per unit of total land used	ECM/Land <sub>total</sub>	t/ha

603

604 **Table 2.** Average milk composition and yield of Langhill dairy production systems605 (mean  $\pm$  s.d.)

Characteristic	Production System <sup>1</sup>			
	LFC	LFS	HFC	HFS
Milk fat (g/kg)	35.4 $\pm$ 0.18	37.8 $\pm$ 0.18	38.2 $\pm$ 0.18	40.0 $\pm$ 0.19
Milk protein (g/kg)	31.5 $\pm$ 0.09	33.7 $\pm$ 0.09	32.1 $\pm$ 0.09	33.5 $\pm$ 0.01
Annual milk yield	9246 $\pm$ 800	10753 $\pm$ 583	7281 $\pm$ 533	8189 $\pm$ 656

<sup>1</sup> LFC = Low Forage Control, LFS = Low Forage Select, HFC = High Forage Control, and LFS = High Forage Select

606

607

608 **Table 3.** Crop yields and on-farm land use by Langhill systems (mean  $\pm$  s.d.)

	Grass silage	Maize silage	Wheat	Pasture
Crop yield (t DM/ha)	10.3 $\pm$ 1.5	11.9 $\pm$ 2.1	11.6 $\pm$ 2.3	10.3 $\pm$ 1.5
Land use by system <sup>1</sup> (ha)				
LFC	18.1 $\pm$ 4.0	4.0 $\pm$ 0.8	3.2 $\pm$ 0.7	
LFS	18.0 $\pm$ 4.2	4.2 $\pm$ 0.7	3.4 $\pm$ 0.9	
HFC	18.2 $\pm$ 4.1	4.1 $\pm$ 0.9	3.3 $\pm$ 0.7	11.9 $\pm$ 2.8
HFS	18.2 $\pm$ 4.2	4.2 $\pm$ 0.8	3.4 $\pm$ 0.8	12.2 $\pm$ 3.0

<sup>1</sup> LFC = Low Forage Control, LFS = Low Forage Select, HFC = High Forage Control, and LFS = High Forage Select. DM = Dry Matter.

609

610

611 **Table 4.** Breakdown of components in purchased feed blends and estimated land

612 use

	Barley	Wheat	Sugar beet pulp	Soya bean meal	Rapeseed meal	Complete blend
Whole crop yield (t DM/ha)	5.9	7.0	10.0	3.0	3.9	
Allocation (%) <sup>1</sup>	100	100	22	80	60	
Percentage in blend (%)						
Low Forage		47	28	25		
High Forage	33.3	33.3			33.3	
Land use (m <sup>2</sup> /kg)						
Low Forage		0.64	0.06	0.75		1.44
High Forage	0.43	0.43			0.70	1.55

<sup>1</sup> Percentage of environmental impact attributed to production of each component of feed blend by mass allocation (Cederberg and Mattsson, 2000).

DM = dry matter.

613

614

615 **Table 5.** Emissions intensity of Langhill dairy production systems expressed as  
 616 kilograms of carbon dioxide equivalents (kgCO<sub>2</sub>e) per functional unit

Functional Unit <sup>2</sup>	Production System <sup>1</sup>				SEM	P-value
	LFC	LFS	HFC	HFS		
LSU (n)	4126 <sup>a</sup>	4398 <sup>ab</sup>	4535 <sup>bc</sup>	4807 <sup>c</sup>	126.3	***
ECM (kg)	0.92 <sup>a</sup>	0.83 <sup>b</sup>	1.10 <sup>c</sup>	1.00 <sup>d</sup>	0.016	***
MS (kg)	12.9 <sup>a</sup>	11.4 <sup>b</sup>	15.2 <sup>c</sup>	13.7 <sup>a</sup>	0.23	***
Land <sub>farm</sub> (ha)	16006 <sup>a</sup>	15971 <sup>a</sup>	11506 <sup>b</sup>	11704 <sup>b</sup>	252.5	***
Land <sub>total</sub> (ha)	6287 <sup>a</sup>	6304 <sup>a</sup>	8041 <sup>b</sup>	7467 <sup>b</sup>	236.2	***
ECM/Land <sub>total</sub> (t/ha)	14.9 <sup>a</sup>	13.4 <sup>a</sup>	21.4 <sup>b</sup>	20.5 <sup>b</sup>	0.82	***

Values within a row with different superscripts differ significantly at  $P < 0.001$

<sup>1</sup>LFC = Low Forage Control, LFS = Low Forage Select, HFC = High Forage Control, and LFS = High Forage Select

<sup>2</sup>Units = livestock units (LSU), total energy corrected milk yield (ECM), total milk solids (MS), on-farm land use (Land<sub>farm</sub>), total land use (Land<sub>total</sub>), and milk yield per unit total land use (ECM/Land<sub>total</sub>)

617

618

619 **Table 6.** Median rankings denoting the relative emissions intensities of Langhill dairy

620 production systems

Functional Unit <sup>2</sup>	Production System <sup>1</sup>				<i>P</i> -value
	LFC	LFS	HFC	HFS	
LSU (n)	1 <sup>a</sup>	2 <sup>b</sup>	3 <sup>b</sup>	4 <sup>c</sup>	*
ECM (kg)	2 <sup>a</sup>	1 <sup>b</sup>	4 <sup>c</sup>	3 <sup>d</sup>	*
MS (kg)	2 <sup>a</sup>	1 <sup>b</sup>	4 <sup>c</sup>	3 <sup>a</sup>	*
Land <sub>farm</sub> (ha)	3 <sup>a</sup>	4 <sup>a</sup>	1 <sup>b</sup>	2 <sup>b</sup>	*
Land <sub>total</sub> (ha)	1 <sup>a</sup>	2 <sup>a</sup>	3 <sup>b</sup>	4 <sup>c</sup>	*
ECM/Land <sub>total</sub> (t/ha)	2 <sup>a</sup>	1 <sup>b</sup>	4 <sup>c</sup>	3 <sup>c</sup>	*

Values within a row with different superscripts differ significantly at  $P < 0.05$ .

<sup>1</sup>LFC = Low Forage Control, LFS = Low Forage Select, HFC = High Forage Control, and LFS = High Forage Select

<sup>2</sup>Units = livestock units (LSU), energy corrected milk yield (ECM), total milk solids (MS), on-farm land use (Land<sub>farm</sub>), total land use (Land<sub>total</sub>), and milk yield per unit total land use (ECM/Land<sub>total</sub>)

621